


IgA nephropathy

Nephropathy - IgA; Berger's disease



http://www.med.niigata-u.ac.jp/npa/Lectures/Images/Slides/IgAN/Exercise/1IgA_PAS1_L.gif

Nephropathy is damage, disease, or other problems with the kidney. IgA nephropathy is a kidney disorder in which antibodies called IgA build up in kidney tissue.

It is also called Berger's disease.

Causes, incidence, and risk factors

IgA is a protein called an antibody that helps the body fight infections. IgA nephropathy (Berger's disease) occurs when too much of this protein is deposited in the kidneys. IgA builds up inside the small blood vessels of the kidney. Structures in the kidney called glomeruli become inflamed and damaged.

IgA nephropathy (Berger's disease) is a form of mesangial proliferative nephritis.

The disorder can appear suddenly (acute), or get worse slowly over many years (chronic glomerulonephritis).

Risk factors include:

- A personal or family history of IgA nephropathy or Henoch Schonlein **purpura**, a form of **vasculitis** that affects many parts of the body
- Caucasian or Asian ethnicity

IgA nephropathy can occur in people of all ages, but it most often affects males in their teens to late 30s.

Symptoms

There may be no symptoms for many years.

Symptoms include:

- Bloody urine that starts during or soon after a respiratory infection
- Repeated episodes of dark or bloody urine
- Swelling of the hands and feet
- Symptoms of **chronic kidney disease**

Signs and tests

IgA nephropathy usually is discovered when a person with no other symptoms of kidney problems has one or more episodes of dark or bloody urine.

There are no specific changes seen during a physical examination. Sometimes, the blood pressure may be high or there may be swelling of the body.

Tests include:

- **Blood urea nitrogen (BUN)** blood test to measure kidney function
- **Creatinine blood test** to measure kidney function
- **Kidney biopsy** to confirm the diagnosis
- **Urinalysis**
- **Urine immunoelectrophoresis**

Treatment

The goal of treatment is to relieve symptoms and prevent or delay **chronic renal failure**. You may get medicines to control high blood pressure and swelling (**edema**), such as angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARBs). Controlling blood pressure is the most important way to delay kidney damage.

Corticosteroids, other drugs that suppress the immune system, and fish oil have also been used to treat this disorder.

Salt and fluids may be restricted to control swelling. A low to moderate **protein** diet may be recommended in some cases.

Some people will need to take medicines to lower their cholesterol.

Eventually, many patients must be treated for chronic kidney disease and may need dialysis.

Expectations (prognosis)

IgA nephropathy gets worse slowly. In many cases, it does not get worse at all. Your condition is more likely to get worse if you have:

- High blood pressure
- Large amounts of **protein in the urine**
- Increased BUN or creatinine levels

About 25% of adults with IgA nephropathy develop **end-stage kidney failure** within 25 years.

Complications

- **Acute nephritic syndrome** or **nephrotic syndrome**
- **Chronic kidney failure**
- **End-stage kidney disease**

Calling your health care provider

Call your health care provider if you have bloody urine or if you are producing less urine than usual.

Reference:

<http://www.ncbi.nlm.nih.gov>

eTOC在校生徒様がeTOCのレッスン
以外の目的で使用することは印刷禁止。
法律で罰せられます。
This document is protected by copyright.
You are breaking the law
if you copy or distribute this file.

