

# Mandibular Dislocation


**Spontaneous mandibular dislocation** usually occurs in people with a history of such dislocations. Although a dislocated mandible is occasionally caused by trauma, the initiating episode is typically a wide opening followed by biting pressure (eg, biting into a large sandwich with hard bread), a wide yawn, or a dental procedure. People prone to dislocation may have naturally loose **temporomandibular joint (TMJ) ligaments**.

Patients present with a wide-open mouth that they are unable to close. Pain is secondary to patients' attempts to close the mouth. If the mandibular midline deviates to one side, the dislocation is unilateral. Although rarely used, a local anesthetic (eg, 2% **lidocaine** 2 to 5 mL) injected into the **ipsilateral** joint and into the adjacent area of insertion of the lateral **pterygoid** muscle may allow the mandible to reduce spontaneously.

Manual reduction may be necessary. Premedication may be used (eg, diazepam 5 to 10 mg IV at 5 mg/min or midazolam 3 to 5 mg IV at 2 mg/min and an opioid such as **mepерidine** 25 mg IV or **fentanyl** 0.5 to 1µg/kg IV) but is usually unnecessary, especially if time will be lost preparing the IV. The longer the mandible is dislocated, the more difficult it is to reduce and the greater the likelihood that dislocation will recur.

Fig. 1

## Mandibular reduction.



The patient's head is stabilized. The operator's thumbs are

placed on the **external oblique** line of the mandible (lateral to the 3rd molar area) or, after wrapping the thumbs in gauze, on the **occlusal surface** of the lower molars. The other fingers are curled under the mandible. The patient is asked to open wide, as if yawning, and the operator then applies downward force on the molars while applying upward force over the chin until the mandible reduces.

Barton bandage may be needed for 2 or 3 days. Most importantly, the patient must avoid opening the mouth wide for at least 6 wk. When anticipating a yawn, the patient should place a fist under the chin to prevent wide opening. Food must be cut into small pieces. If the patient suffers from chronic dislocations and more conservative treatment modalities have been exhausted, an oral and **maxillofacial surgeon** may be consulted. As last-resort treatments, the ligaments around the TMJ can be surgically tightened (shortened) in an attempt to stabilize the joint or the **articular eminence** can be reduced (**eminectomy**).

eTOC在校生徒様がeTOCのロゴを  
以外の目的で使用すること及び印刷  
法律で罰せられます。  
This document is protected by copyright.  
You are breaking the law  
if you copy or distribute this file.

**eTOC**  
English Teachers On Call